

SAFETY

Wear suitable gloves, overalls and face shield when handling this product.

STORAGE

Store away from direct sunlight and in frost free conditions, preferably between 5°C and 35°C. Crystallisation can be a product characteristic under extreme conditions.

DISPOSAL

Do not allow product to enter water courses un-diluted.

Triple rinse emptied containers, adding the rinsate to the spray tank. Handle empty containers and their residues as established and in accordance with local legislation and guidance.

WARRANTY

Safagrow guarantees the content of the components as indicated on the label. The buyer assumes all risks regarding the use and management.

INSTRUCTIONS FOR USE

Read the label before use.

Shake the bottle well before use.

Half fill spray tank with water and begin agitation. Empty the required quantity into the spray tank, mix until dissolved or dispersed and add the balance of the water. Ensure agitation is maintained throughout the spraying operation. Do not allow the spray tank agitation to stop whilst product remains in the tank.

COMPATABILITY

Max P is compatible and can be mixed with many pesticides and nutrients but not all. However, Safagrow cannot accept any liability for any loss or damage resulting from tank mixes, as it is impossible to screen all possible mixes and conditions.

Safagrow recommends the buyer tests the proposed mix on small scale first.

H412:	Harmful to aquatic life with long lasting effects.
P273:	Avoid release to the environment.
P501:	Dispose of contents/container to hazardous or special waste collection point.

APPLICATION RECOMMENDATIONS

CROP	RATE	TIMING
Cereals	2.0 ml/l	Apply early in the crop cycle
Vegetables	2.0 ml/l	Apply 20 days after transplant and then repeat 14 days later
Cabbage, Cauliflower	2.0 ml/l	Apply 20 days after transplant
Peppers, Tomatoes, Brinjals	2.0 ml/l	Apply 20 days after transplant and then repeat 14 days later
Ornamentals, Greenhouse, Polyhouse	2.0 ml/l	Apply early in the growth period and repeat after 14 days
Other Crops	2.5 ml/l	Where Phosphorous deficiency is observed apply and repeat after 14 days
Additional Crops: For additional crops not listed here, check our website at WWW.SAFAGROW.COM		

EC FERTILISER

NPK – fertiliser suspension 6.3 – 25.2 – 12.6 with micro-nutrients

Nitrogen (N) Total	6.3%w/w
4.1%w/w Ammoniacal nitrogen	
2.2%w/w Ureic nitrogen	
Phosphorus Pentoxide (P ₂ O ₅) Water-soluble	25.2%w/w
Potassium Oxide (K ₂ O) Water-soluble	12.6%w/w
Boron (B) Water-soluble	0.022%w/w
Copper (Cu) Water-soluble	0.054%w/w
Iron (Fe) Water-soluble	0.110%w/w
0.110%w/w chelated by EDTA	
Manganese (Mn) Water-soluble	0.054%w/w
Molybdenum (Mo) Water-soluble	0.001%w/w
Zinc (Zn) Water-soluble	0.054%w/w
Chelated fraction stable within pH range 3–10	