

# Calcium Aloy®



## SAFETY

Wear suitable gloves, overalls and face shield when handling this product.

#### STORAGE

Store away from direct sunlight and in frost free conditions, preferably between 5°C and 35°C. Slight thickening can be a product characteristic under extreme conditions.

# DISPOSAL

Do not allow product to enter water courses un-diluted.

Triple rinse emptied containers, adding the rinsate to the spray tank. Handle empty containers and their residues as established and in accordance with local legislation and guidance.

#### WARRANTY

Safagrow guarantees the content of the components as indicated on the label. The buyer assumes all risks regarding the use and management.

### COMPATABILITY

Calcium Aloy® is compatible and can be mixed with many pesticides and nutrients but not all. However, Safagrow cannot accept any liability for any loss or damage resulting from tank mixes, as it is impossible to screen all possible mixes and conditions.

Safagrow recommends the buyer tests the proposed mix on small scale first.



# WARNING

H319:	Causes serious eye irritation.
H412:	Harmful to aquatic life with long lasting effects.
P264:	Wash hands thoroughly after handling.
P273:	Avoid release to the environment.
P280:	Wear protective gloves/protective clothing/eye protection/ face protection.
P305+351+338:	IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
P337+313:	If eye irritation persists: Get medical attention.
P501:	Dispose of contents/container to hazardous or special waste collection point.

VAL1565BUK

# INSTRUCTIONS FOR USE

dose rates.

Always shake the bottle well before use.

NEVER MIX PESTICIDES IN THEIR CONCENTRATED FORM.

If Spray volumes are less than 200I/ha or if you are in doubt, then perform a "Jar test" on the mix before starting.

Half fill the tank with water and switch on agitation. Always make sure the additions are fully mixed until dissolved or dispersed before adding the next product.

- Add water conditioners, if needed. Add granules powders and mix well.
   Pre disperse if possible (WP. DG).
- Add Calcium Alov® and other flowables (SC, SE).
- Add Emulsion Concentrates and crop oils (EC).
- Add water soluble: liquids and solids, (SL, SP).
- . Add wetters, if needed and top up with water,

Provided these guidelines are followed almost all tank mixes are physically compatible, with continuous agitation. If agitation fails, for any reason, then it is essential to clean out the tank immediately.

#### APPLICATION RECOMMENDATIONS CROP RATE TIMING Apply the product in a minimum of 200 litre of water unless specified Apply and repeat at 10 day intervals from 4 weeks Apples and Pears 3-5I/ha after petal fall. Water rate 500 L/Ha minimum. Broccoli, Cabbage, Calabrese Apply from 4-9 leaves and repeat at 10-14 day 3-5I/ha Cauliflower and intervals. Sprouts In case of moderate to severe deficiency repeat same Carrots 3-51/ha rate at 10 to 14 day intervals. Apply and repeat at 7 day intervals during fruit Cucurbits 3-51/ha (field grown) development. Apply 10-14 days from transplanting or emergence Lettuce 3-5I/ha (field grown) and repeat as required in intervals of 10-14 days. Onions Apply 10-14 days from transplanting or emergence 3-51/ha (field grown) and repeat as required in intervals of 10-14 days. Apply 14 days after emergence and repeat as Potatoes 5I/ha necessary at the same rate in intervals of 10-14 days.

Additional Crops: For additional crops not listed here, check our website at WWW.SAFAGROW.COM

EC FERTILISER		
Micro-nutrient fertiliser		
Manganese (Mn) Total	1.5%w/w	
1.5%w/w Nitrate		
Zinc (Zn) Total	0.6%w/w	
0.6%w/w Oxide		
To be used only where there is a recognised need. Do not exceed the appropriate		